

# UNCLASSIFIED

AD NUMBER
AD291735
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; 10 OCT 1962. Other requests shall be referred to Army Medical Research Laboratory, Fort Knox, KY.
AUTHORITY
usamrl ltr, 26 feb 1970

THIS PAGE IS UNCLASSIFIED

**UNCLASSIFIED**

**AD 291 735**

*Reproduced  
by the*

**ARMED SERVICES TECHNICAL INFORMATION AGENCY  
ARLINGTON HALL STATION  
ARLINGTON 12, VIRGINIA**



**UNCLASSIFIED**

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

291735

# US ARMY MEDICAL RESEARCH LABORATORY

FORT KNOX, KENTUCKY

REPORT NO. 554


THE SYNERGISM OF AUTONOMIC DRUGS ON OPIATE OR  
OPIOID-INDUCED ANALGESIA: A DISCUSSION OF ITS  
POTENTIAL UTILITY AND AN ANNOTATED  
BIBLIOGRAPHY

Capt Wayne O. Evans, MSC

CATALOGED BY ADIA  
AS AD NO.

291735

AST  
DEC 27 1962  
TISIA C



UNITED STATES ARMY  
MEDICAL RESEARCH AND DEVELOPMENT COMMAND 10 October 1962

NO OTS

Report Submitted 29 August 1962

Author

Capt Wayne O. Evans, MSC  
(Ph. D.)

Psychomotor Branch  
Psychology Division

Qualified requestors may obtain copies of this report from ASTIA.

Foreign announcement and dissemination of this report by ASTIA  
is limited.

REPORT NO. 554

**THE SYNERGISM OF AUTONOMIC DRUGS ON OPIATE OR  
OPIOID-INDUCED ANALGESIA: A DISCUSSION OF ITS  
POTENTIAL UTILITY AND AN ANNOTATED  
BIBLIOGRAPHY**

**Capt Wayne O. Evans, MSC**

**Psychology Division  
US ARMY MEDICAL RESEARCH LABORATORY  
Fort Knox, Kentucky**

**10 October 1962**

**Pharmacology of the Combat Soldier  
Task 04  
Internal Medicine  
USAMRL Project No. 6X60-01-001**

Report No. 554  
USAMRL Project No. 6X60-01-001-04

### ABSTRACT

## THE SYNERGISM OF AUTONOMIC DRUGS ON OPIATE OR OPIOID-INDUCED ANALGESIA: A DISCUSSION OF ITS POTENTIAL UTILITY AND AN ANNOTATED BIBLIOGRAPHY

### OBJECT

The object of this paper is to gather together information pertaining to the potentiation of opiate-induced analgesia by autonomically active drugs to facilitate theoretical explanations of their action and considerations of the practical uses of this type of mixture.

### RESULTS

The evidence indicates that combining an opiate with any one of a diverse group of autonomic drugs will result in an increase in the degree of induced analgesia. D-amphetamine was particularly considered since its addition to an opiate increases the analgetic effect by 60 to 100%. At the same time, d-amphetamine helps minimize the undesirable side effects of opiates by reducing the degree of nausea, constipation, sedation and mental depression; and by normalizing blood pressure, oxygen consumption, and respiratory rate.

### CONCLUSION

The combination of d-amphetamine with an opiate for the induction of analgesia should offer certain unique advantages for use in civil disasters, combat military medicine, and in other similar situations in which environmental exposure, hemorrhage, and fear contribute to a danger of death due to shock. Its use would also include situations in which sedation is contraindicated due to a need to keep the patient ambulatory and cooperative.

### RECOMMENDATION

The possibility of combat military use of this mixture should be given serious consideration. Studies should be conducted to provide an explanation for the potentiation effect and to seek further evidence of its practical applicability.

APPROVED:

George S. Harker  
GEORGE S. HARKER, Ph. D.  
Director, Psychology Division

APPROVED:

Floyd A. Odell  
FLOYD A. ODELL, Ph. D.  
Technical Director of Research

APPROVED:

Sven A. Bach  
SVEN A. BACH  
Colonel, MC  
Commanding



# THE SYNERGISM OF AUTONOMIC DRUGS ON OPIATE OR OPIOID-INDUCED ANALGESIA: A DISCUSSION OF ITS POTENTIAL UTILITY AND AN ANNOTATED BIBLIOGRAPHY

## I. INTRODUCTION

Two major considerations have tended to restrict the utility of opiate and opioid alkaloids as analgesic agents: first, their tendency to produce tolerance and addiction; and second, the problems introduced by their side effects of respiratory depression, reduced blood pressure, mental depression, constipation, nausea, and sedation.

The problem of tolerance and addiction has not too greatly affected combat military usage since, generally, a trauma case receives opiates for a comparatively short period of time. However, the side effects produced by the actions of opiates on the autonomic nervous system do constitute a problem, especially in those situations in which shock is present or imminent. Shock is a very common danger when injury occurs under exposed environmental conditions, and the side effects of the opiates are such as to intensify shock, or to increase the likelihood of the appearance of profound shock (1, 2).

Considering these difficulties inherent in the use of opiates, a method of increasing their analgetic activity while reducing their depressant and shock enhancing effects would be of great use in situations in which shock, ability to carry out instructions, or the necessity of maintaining consciousness, contraindicates the use of an opiate alone as an analgesic.

## II. DISCUSSION

A rather diverse group of autonomically active agents have been shown to potentiate the analgetic activity of opiates. The majority of the investigations relating to this effect are reported in the accompanying annotated bibliography. Evidence for potentiation has been put forth for parasympathomimetic agents such as neostigmine (3) and physostigmine (4), parasympatholytic drugs such as atropine and scopolamine (5), sympathomimetic agents such as d-amphetamine (6) and methamphetamine (7), sympatholytic drugs like chlorpromazine (8) and dibenzylamine (9), and drugs of the tryptamine and tryptophane type (10). Furthermore, there is no evidence of a relationship between the particular physiological changes usually considered in connection with

these drugs and the degree to which they are able to potentiate opiates (11).

At the present time we have no adequate theoretical explanation of this potentiation. Investigators have suggested three theories of action: 1) that an excitation of the central sympathetic nervous system produces analgesia and that the morphine effect is due to a release of adrenaline which facilitates sympathetic action (12), 2) that, since opiates act as cholinesterase inhibitors and morphine is potentiated by cholinesterase inhibitors, the induced analgesia is a "cholinergic" event (13); and 3) that, since all of the agents which potentiate morphine can be structurally related to the hypothesized central adrenergic "alpha" and "beta" receptors, the action of potentiation is through an active competition with morphine for adrenergic sites, thus leaving more opiate available for analgetic action (9). At present, the evidence available discounts the first two theories (5, 11, 14, 10) and not enough evidence exists to support the "adrenergic competition" hypothesis. However, regardless of the present lack of a theoretical explanation of this potentiation, empirical evidence does demonstrate the fact of its existence and its potential usefulness in combat military medicine or situations of civil disaster.

Of the various drugs which will potentiate an opiate-induced analgesia, d-amphetamine seems to offer the greatest practical utility because it counteracts the shock-enhancing aspects of morphine action. Also, more evidence exists as to its clinical effects when combined with opiates than for the other agents. The first mention of a true potentiation of an opiate-induced analgesia by d-amphetamine was presented by Ivy and his colleagues in 1944 (6, 15, 16). However, as early as 1941, Abreu and Handley (17) had shown that amphetamine could normalize respiratory depression and oxygen consumption in morphinized rats. Similarly, in 1941, DeVoine Guyot (18) reported that patients with coronary occlusion when treated with 0.5 gr to 0.75 gr of morphine alone would show more vomiting, inhibition of bowel action, respiratory depression, hypotension, and mental depression than when the same dose of morphine was given in combination with 10.0 mg of r-amphetamine. Also, Morrison and Abreu (19) showed that the depression of oxygen uptake in dogs by 10 mg/kg of morphine could be combated by 0.5 mg/kg of l-amphetamine, d-amphetamine, or r-amphetamine. In this study the d-amphetamine raised the depressed oxygen consumption by 22% and was superior to the racemic or levo-rotary forms.

In 1944, Ivy and his colleagues demonstrated that d-amphetamine could increase an opiate-induced analgesia in dogs (15), in mice (16),

and in man (6). In man, 16 mg of morphine sulfate in combination with 20 mg of d-amphetamine sulfate produced a degree of analgesia 60% greater than with the same dose of morphine alone. At the same time, the mixture overcame the depression of morphine on respiration and blood pressure, reduced nausea and vomiting, and normalized the critical flicker fusion threshold and choice reaction time of the subjects.

Between 1944 and 1950 the potentiation of opiate and analgesia by d-amphetamine was confirmed by Nickerson and Goodman (20) with normal subjects given isonipecaine. Also, Abel and Harris (21) showed the effect with obstetrical patients. During the same period, Handley and his colleagues (22, 23) reconfirmed the normalizing effects of amphetamine on respiration, pulse rate, and blood pressure of morphinized patients. They also showed that d-amphetamine was superior to metrazol, ephedrine, caffeine, and nikethamide in combating opiate side effects. Nickerson (24) in 1950 by testing the reduction of pain induced by immersion of the hand in ice water showed the opiate potentiating effect of d-amphetamine in humans by combining meperidine (100 mg) and d-amphetamine (10 mg).

The first large scale clinical test of an opiate mixed with d-amphetamine was conducted in 1951 (25). Abel and his associates demonstrated with 7,000 obstetrical cases that adding 5 mg of d-amphetamine to 10 mg of morphine would provide good analgesia with the minimum of side effects and also shorten the length of time to the first inspiration by newborn infant as compared to the time of inspiration of the infant for patients given morphine alone.

Since 1951 only a few investigators have shown interest in this problem area. Saxena and Gupta (26), Guseva (27), Matsumura et al (28), Witkin et al (29), and Evans (9, 30), have reconfirmed in laboratory studies the potentiation actions of amphetamine on opiate analgesia and sought theoretical explanations for the effect, but clinical interest seems to have waned.

Perhaps the explanation of the decline of clinical interest lies in the fact that in a modern hospital shock does not present the major problem that it once did. Thus, there is no pressing requirement for a morphine compound with little or no shock enhancing property even if it allows a reduction in the amount of opiate needed to obtain a satisfactory level of analgesia. Similarly, the overcoming of the mental depression and sedation produced by morphine has little advantage in the modern hospital.

On the other hand, the use of morphine in a civil disaster or in the combat military situation may be quite different in terms of the requirements placed upon the drug. The administration of 15 mg morphine from a pre-packaged syringe to a wounded man in a wet, muddy foxhole during a Korean winter presents problems not present in a hospital in the United States. In combat or in situations of civil disaster, every condition is present to maximize the possibility of death due to shock. It is in situations of this nature that the full potential of the mixture of an opiate with d-amphetamine would be realized.

### III. CONCLUSION

It has been demonstrated in dogs, rats, mice, healthy humans, and patients that the addition of d-amphetamine to an opiate will increase its analgetic potency from 60 to 100%. At the same time the addition of d-amphetamine helps normalize the opiate side effects of respiratory depression, lowered oxygen consumption, hypotension, mental depression, constipation, and emesis.

It would seem that this combination could be of great use as a combat military analgesic and its potential should be further studied.

### IV. REFERENCES

1. Editorial: The (Ab) use of morphine. U.S. Armed Forces Med. J. 1: 247, 1950.
2. Jeghers, H. and J. B. Brick. Hazards in the therapeutic use of morphine. In: Medical Clinics of North America, W. B. Saunders, Co., Philadelphia, 1950.
3. Slaughter, D. Neostigmine and opiate analgesia. Arch. Int. Pharmacodyn. 83: 143, 1950.
4. Porszasz, J., J. Knoll and E. Komlos. Wirkung der Parasympathomimetika auf die analgesia. Acta Physiol. Acad. Sci. Hung. 2: 469, 1951.
5. Knoll, J. and E. Komlos. Die Analgetische Wirkung des Atropins und sein Synergismus mit Morphin und Prostigmin. Acta Physiol. Hung. 2: 57, 1951.
6. Ivy, A. C., F. R. Goetzl and D. Y. Burrill. Morphine-dextroamphetamine analgesia. War Med. 6: 67, 1944.

7. Milosevic, M. P. The analgesic activity of sympathomimetic amines. *Acta Med. Jugosl.* 12: 111, 1958.
8. Courvoisier, S., J. Fournel, R. Ducrot, M. Kolsky and P. Koetschet. Propriétés pharmacodynamique du chlorhydrate de chloro-3-(diéthylamine-3-propyl)-10-phénothiazine (4560 RP), étude expérimentale d'un nouveau corp utilisé dans l'anesthésie potentialisée et dans l'hibernation artificielle. *Arch. Int. Pharmacodyn.* 92: 305, 1953.
9. Evans, W. O. The potentiation of opiate induced analgesia by stimulant drugs. II: The effect of sympathomimetics. USAMPL Report, Ft Knox, Ky (in preparation).
10. Sigg, E. B., G. Caprio and J. A. Schneider. Synergism of amines and antagonism of reserpine to morphine analgesia. *Proc. Soc. Exper. Biol. and Med.* 97: 97, 1958.
11. Knoll, J., E. Komlos and J. Porszasz. Analgesie und Cholinesterasehemmung. *Acta Physiol. Hung.* 2: 478, 1951.
12. Gross, E. G., Helen Holland, H. R. Carter and Eunice Christensen. The role of epinephrine in analgesia. *Anesthesiol.* 9: 459, 1948.
13. May, E. L. Analgesics. In: *Medicinal Chemistry*, Alfred Berger (Ed.), Interscience Publishers, Inc., New York, 1960.
14. Schaumann, W. Zur Hypothese eines cholinergen Wirkungsmechanismus des Morphins. *Naunyn Schmiedeberg's Arch. Exp. Path. Pharmacol.* 237: 229, 1959.
15. Goetzl, F. R., D. Y. Burrill and A. C. Ivy. The analgesic effects of sympathomimetics and the combination of morphine with amphetamine. *Proc. Inst. Med. Chicago*, 15: 87, 1944.
16. Goetzl, F. R., D. Y. Burrill and A. C. Ivy. The analgesic effect of morphine alone and in combination with dextro-amphetamine. *Proc. Soc. Exper. Biol. and Med.* 55: 248, 1944.
17. Abreu, B. E. and C. A. Handley. The effects of benzedrine on respiration and oxygen consumption in men and rats. *U. Calif. Pub. Pharmacol.* 2: 99, 1941.

18. DeVoine Guyot, J. The use of benzedrine sulfate to overcome the untoward effects of morphine in the treatment of coronary occlusion. J. Mo. Med. Assn. p. 93, March 1941.
19. Morrison, J. L. and B. E. Abreu. Effects of d-, l-, and r-amphetamine on oxygen consumption of morphinized dogs. Fed. Proc. 2: 88, 1943.
20. Nickerson, M. and L. S. Goodman. Synergistic isonipecaine-amphetamine analgesia. Fed. Proc. 6: 360, 1947.
21. Abel, S. and S. C. Harris. Morphine-benzedrine analgesia in obstetrics. (Abstract) Amer. Physiol. Soc. Proc. p. 67, 1947.
22. Handley, C. A., D. L. Ensberg and H. M. Sweeney. The relative efficiency of central nervous system stimulants in respiratory depression from morphine. Fed. Proc. 3: 75, 1944.
23. Handley, C. A. and D. L. Ensberg. A comparison of amphetamine sulfate with other stimulants of the central nervous system in morphine respiratory depression. Anesthesiol. 6: 561, 1945.
24. Nickerson, M. Analgesic effects of nitrous oxide and meperidine alone and combined with amphetamine. West. Soc. Clin. Rsch. p. 541, April 1950.
25. Abel, S., Zelda B. Ball and S. C. Harris. The advantages to mother and infant of amphetamine in obstetrical analgesia. Amer. J. Obst. Gynecol. 62: 15, 1951.
26. Saxena, P. N. and G. P. Gupta. Analgesic potentiating effects of ephedrine and methamphetamine. J. Indian Med. Prof. 4: 1553 and 1599, 1957.
27. Guseva, E. N. Combined action of analgetic morphine substitutes with analeptics. Farmakol. I. Toksikol. 20: Supl. 1, 1957.

28. Matsumura, M., S. Takaori and R. Inaki. Suppressive synergism of morphine with methamphetamine on the afferent pathways of the central nervous system in the cat. *Jap. J. Pharmacol.* 9: 67, 1959.
29. Witkin, L. B., M. Maggio, E. O'Keefe and F. Galdi. A study of some central nervous stimulants in mice. *Fed. Proc.* 19: 269, 1960.
30. Evans, W. O. The potentiation of opiate-induced analgesia by stimulant drugs. I: The effect of monoamine oxidase inhibitors and caffeine. USAMRL Report No. 519, Ft Knox, Ky., 1961.

AN ANNOTATED BIBLIOGRAPHY OF THE SYNERGISM OF  
ANALGESIA OF OPIATES BY AUTONOMIC DRUGS

ABEL, S., ZEIDA B. BALL and S. C. HARRIS

The advantages to mother and infant of amphetamine in obstetrical analgesia. Amer. J. Obst. and Gynecol. 62: 15-27, 1951.

A clinical study of 7,000 cases given 1/6 gr of morphine alone or with 5 mg of d-amphetamine. Also 100 mg of demerol alone and with 1/150 gr of scopolamine. Clinical estimates of morphine-amphetamine on degree of analgesia were favorable. The first inspiration of infant was significantly retarded by morphine alone and normalized by the morphine-amphetamine mixture.

ABEL, S. and S. C. HARRIS

Morphine-benzedrine analgesic in obstetrics. (Abstract) Paper at Amer. Physiol. Soc. Proceedings, p. 67, 1947.

Nine patients were given 1/4 gr of morphine plus 5 mg of benzedrine for labor pain. First inspiration of infants born one to seven hours after drug was 42 sec mean (0-97). Good analgesia during labor was obtained as well as reducing the time to first inspiration by the infant.

ANGIBEAUD, P., L. BUCHEL and J. LEVY

Sur quelques phénomènes de synergie. I. Associations d'analgésiques et d'un spasmolytique considéré comme leur analogue structural. C. R. Soc. Biol. (Paris), 149: 258-261, 1955.

Using mice with a method of thermal and mechanical stimulation, the analgetic action of 1-amidone, pethidine and morphine was measured alone and in combination with 2-(1-piperidyl) ethyl ester hydrochloride of cyclohexylcyclohexanecarboxylic acid. Synergism was shown with 1-amidone, but little effect with pethidine and none with morphine.

ABREU, B. E. and C. A. HANDLEY

The effects of benzedrine on respiration and oxygen consumption after morphine in men and rats. U. Calif. Pub. Pharmacol. 2: 99-104, 1942.



Six healthy men received 0.3 mg/kg morphine  $\text{SO}_4$  orally. They were measured one hour later for:  $\text{O}_2$  consumption, respiration, cardiac rate, pain by force of a dull point into hand. Also,  $\text{O}_2$  from rats after 20 mg/kg morphine + 10 mg/kg of benzedrine. 0.3 mg/kg benzedrine used in humans.

0.3 mg/kg of morphine did not significantly depress  $\text{O}_2$  consumption in humans nor could benzedrine be shown to have any effect to antagonize. In rats, morphine depression and benzedrine antagonism were shown.

**BOREUS, L. O. and F. SANDBERG**

The influence of three phenothiazine derivatives and of amphenazole on the action of methadon. Studies with two algometric methods in untrained human subjects. *J. Pharm. (London)*, 11: 449-455, 1959.

Chlorpromazine and acepromazine showed some analgesic activity using heat to the thumb nail as a test. Mepazine had no activity. By this test chlorpromazine and acepromazine did not potentiate methadone and mepazine antagonized its analgetic action. Amphenazole had no effect on methadone and a slight activity by itself.

**BUCHÉL, L., J. LÉVY and O. TANGUY**

Sur la potentialisation de l'action analgesique de la 1-méthadone par la 5-hydroxytryptamine (sérotonine). *C. R. Acad. Sci.* 246: 2947-2949, 1958.

Serotonin enhances analgetic action of 1-methadon using mechanical stimuli. Reserpine promoted release of serotonin but antagonized methadon.

**CHANG TAN-MU, TAH-CHAO FONG and FU-HAN LUE**

Potentiating effects of diphenhydramine on the analgesic action of Han-Fang-Chi and some other analgesics. *Acta Physiol. Sinica*, 21: 133-141, 1957.

Using the method of Reinhard et al the alkaloids of *Stephania tetrandra* S. Moore (tetrandrine) and also morphine, pethidine and phenazone were shown to be analgetic alone. Diphenhydramine

significantly potentiated the analgetic action of all of these compounds.

CHEN, J. Y. P.

Analgesic-potentiating and diuretic effects of 1-dl-methylamino-3-phenyl-4-methyl-hexane HCl (z-4) and 1-dimethylamino-2-phenyl-3-methyl-pentane HCl (z-134). J. Pharmacol. exp. therap. 117: 451-460, 1956.

Used hot plate with mice and found a true potentiation of 7 mg/kg of morphine by: z-134, 20 mg/kg; z-4, 20 mg/kg; chlorpromazine, 2 mg/kg. Similar results with demerol and tail pinching method. Drugs not analgetic at these doses by themselves. z-4 and z-134 were respiratory stimulants in pentobarbitalized dogs. Also, lowered intestinal tone. z-4 and z-134 increased urine output by 500% over saline controls. Both produce a transitory fall in blood pressure. No hypnotic potentiating effects with pentobarbital. Toxic dose of 93 mg in rat produces central excitement and convulsions.

CHRISTENSEN, E. M. and E. G. GROSS

A comparison of the analgesic effects on human subjects of 6 dimethylamino-4-4 diphenyl-3-heptanone (AN148), morphine and meperidine (demerol) and the relative efficiency of AN148 for preoperative and postoperative use. J.A.M.A. 137: 594-599, 1948.

Used Hardy, Wolff, and Goodell method with eleven trained human subjects. Subjects guessed compounds used sometimes. Gave either 0.3 mg of atropine or scopolamine with methadon (2.5 mg), morphine (10 mg) and meperidine (50 mg). In all cases period of analgesia was shortened and reduced threshold increases of analgesic. Neostigmine at a dose of 0.5 mg caused an increase in threshold by itself and also a synergism with opiates.

CHRISTIE, G., S. GERSHON, R. GRAY, F. H. SHAW, I. McCANCE and D. W. BRUCE

Treatment of certain side effects of morphine. Brit. Med. J. 5072: 675-680, 1958.

Normal humans given 20 mg or 30 mg of morphine alone or with amiphenazole or cyclizine. Both reduced vomiting, nausea or sleepiness of morphine. Amiphenazole at 40-100 mg particularly helpful. No respiratory depression.

COOK, L., GERALDINE NAVIS and E. J. FELLOWS

Enhancement of the action of certain analgetic drugs by  $\beta$ -diethyl-aminoethyldiphenylpropylacetate hydrochloride (SKF No. 525A). J. Pharmacol. exp. therap. 112: 473-479, 1954.

The investigators used tail flick method in rats and found a true potentiation at 100 mg/kg of SKF 525A of morphine; 2 mg of 4.6x, meperidine, 10 mg of 2.2x; also of demerol, 3 mg, methorphan 0.3 mg and codeine 60 mg. Did not effect morphine induced depression of respiration. SKF 525A enhances analgetic response of morphine tolerant rats.

COURVOISIER, S., J. FOURNEL, R. DUCROT, M. KOLSKY and P. KOETSCHET

Propriétés pharmacodynamique du chlorhydrate de chloro-3 (diméthylamine-3-propyl)-10-phénothiazine (4560RP), étude expérimentale d'un nouveau corp utilisé dans l'anesthésie potentialisée et dans l'hibernation artificielle. Arch. internat. pharmacodyn. 92: 305-561, 1953.

The investigators used the hot plate method with mice and found that chlorpromazine produced marked increases in analgetic potency and time of action of morphine, meperidine, aspirin, salicylamide, aminopyrine and phenacetin.

DE JONGH, D. K.

Remarks on the mechanism of analgesic action of morphine. Acta Physiol. Pharmacol. Neerl. 3: 164-172, 1954.

Using radiant heat stimulus in guinea pigs (method of Winder), tested morphine (4 mg/kg), salicylamide (50 mg/kg), prostigmine (0.025 mg/kg), atropine (1.0 mg/kg) alone and in combination. Each drug by itself raised the threshold. Morphine plus prostigmine was not even additive while atropine plus morphine was additive. Prostigmine and atropine were antagonistic. The results do not suggest that morphine acts through a cholinergic mechanism.

**DE VOINE GUYOT, J.**

The use of benzedrine sulfate to overcome the untoward effects of morphine in the treatment of coronary occlusion. J. Mo. St. Med. Assn. pp. 93-94, March 1941.

Human patients with coronary occlusion treated with 1/2 to 3/4 gr morphine plus 10 mg of benzedrine orally. Vomiting was controlled, bowel action inhibited far less, the blood pressure fall of 50 to 75 mm with morphine alone was reduced to 20 mm by combination. Mental depression was minimized.

**EVANS, W. O.**

The potentiation of opiate induced analgesia by stimulant drugs.  
I: The effect of monoamine oxidase inhibitors and caffeine.  
USAMRL Report No. 519, Ft Knox, Ky., 1961.

Using the jump-flinch method in rats codeine  $\text{SO}_4$  was found at a dose of 30 mg/kg to be potentiated by about 65% by d-amphetamine (2.5 mg/kg) and antagonism by about 40% by caffeine citrate (12 mg/kg). Iproniazid (40 mg/kg),  $\beta$ -phenylisobutylhydrazine (5.0 mg/kg) and  $\beta$ -phenylisopropylhydrazine (5 mg/kg) were found to have no effect.

**EVANS, W. O.**

The potentiation of opiate induced analgesia by stimulant drugs.  
II: The effect of sympathomimetics. USAMRL Report, Ft Knox, Ky. (in preparation).

Using the jump-flinch method in rats 16 mg/kg of morphine  $\text{SO}_4$  was found to be potentiated by 100% by both d-amphetamine (5 mg/kg) and methamphetamine (5 mg/kg). Morphine analgesia was also potentiated by about 60% by trimethylamine HCl (30 mg/kg), 3,4-dihydroxybenzaldehyde (20 mg/kg) and 3,4-dichlorobenzaldehyde (20 mg/kg). Isopropylamine (2 mg/kg), n-propylamine (2 mg/kg) and sec-butylamine (5 mg/kg) were found to antagonize morphine by about 50%. Pre-treatment with 30 mg/kg of phenoxybenzamine was found to potentiate morphine analgesia. The results suggest a relationship to alpha and beta adrenergic receptors, perhaps a competition of adrenergic agents and morphine.

**FLODMARK, S. and T. WRAMNER**

The analgetic action of morphine, esserine and prostigmine studied by a modified Hardy-Wolff-Goodell method. *Acta Physiol. Scand.* 9: 88-96, 1945.

Using 19 human subjects with radiant heat method to compare 15 mg of morphine with 3 mg of morphine plus 0.5 mg of prostigmine. Mixture was 40% more analgetic. Prostigmine at 1 mg and also esserine at 1 mg produced slight rises in threshold of pain.

**GERSHON, S., D. W. BRUCE, N. ORCHARD and F. H. SHAW**  
Amiphenazale and morphine in the production of analgesia. *Brit. Med. J.* 5092: 366-368, 1958.

500 cases treated with morphine and amiphenazale (2-4, diamino-5-phenylthiazole) (in doses up to 200 mg t.i.d.). Morphine may be given in large doses without risk. No sedation found, addiction to morphine doesn't develop with months of treatment. No euphoria is present with mixture. No withdrawal symptoms.

**GOETZL, F. R., D. Y. BURRILL and A. C. IVY**

The analgesic effect of morphine alone and in combination with dextroamphetamine. *Proc. Soc. Exper. Biol. and Med.* 55: 248-250, 1944.

Using mice with tail pinching method given 20 mg/kg of morphine  $SO_2$  and 35 mg/kg of d-amphetamine. Mixture produced an increase in threshold to squeal of about 50% and eliminated the usual morphine induced Straub reaction. Action developed within 60 min to max and lasted six hours to base line return.

**GOETZL, F. R., D. Y. BURRILL and A. C. IVY**

The analgesic effects of sympathomimetics and the combination of morphine with amphetamine. *Proc. Inst. Med. Chicago*, 15: 87-88, 1944.

Using voltage to metal tooth filling in dogs and man. Used first twitch or first pain report. Found d-amphetamine enhanced analgesia. Results confirmed with tail pressure to squeal in mice.

GRIMMETT, M. R., K. D. NEAME and F. N. FASTIER

Interaction between 4-methyl-2 amino-pyridine (W45) and certain other drugs, particularly morphine. Proc. Univ. Otago. Med. Sch. 38: 9-10, 1960.

W45 has some analgetic potency of its own and also enhances that of morphine using pricking of a rat's ear as the method in doses 20-40 mg/kg of morphine and 2.0 to 6.0 of W45. Its own analgetic effects are not antagonized by nalorphine. Its own effect was increased by a pre-treatment with phenelzine but not effected by reserpine or amphetamine.

GRODEN, B. M.

Daptazole: an aid to morphine administration. Surgo, (Glasgow), 23: 122-125, 1957.

A review of the literature on the combination of amiphenazole and morphine. Concludes the combination is clinically useful.

GUSEVA, E. N.

Combined action of analgetic morphine substitutes with analeptics. Farmakol. I. Toksikol. 20: Suppl. 1, 1957 (Russian Text).

Morphine and morphine like drugs were combined with amphetamine (30 mg/kg), caffeine (0.3 mg/kg) or 5 (2-bromoallyl)-5-isopropylbarbituric acid (30 mg/kg). Mice were used to test analgetic potency. All potentiated morphine. Larger doses of caffeine or amphetamine antagonized morphine.

HANDLEY, C. A.

A comparison of the action of amphetamine with other central nervous system stimulants in morphine respiratory depression. Proc. and Trans. Texas Acad. Sci. 28: 95, 1945.

Same material as in Anesthesiology, 6: 561-564, 1945.

HANDLEY, C. A. and D. L. ENSBERG

A comparison of amphetamine sulfate with other stimulants of the central nervous system in morphine respiration depression. Anesthesiology, 6: 561-564, 1945.

Fourteen humans took 0.5 mg/kg s. c. of morphine SO<sub>4</sub>. Five studied with each drug.

	Dose in mg/kg	Minute Vol. % increase over depressed state
Amphetamine SO <sub>4</sub>	0.2	102
Caffeine and Na Benzoate	10.0	17
Ephedrine SO <sub>4</sub>	0.5	30
Metrazol	5.0	21
Nikethamide	7.5	37

Amphetamine and ephedrine also raised morphine depressed pulse and blood pressure. None of the others did this.

HANDLEY, C. A., D. ENSBERG and H. M. SWEENEY

The relative efficiency of central nervous system stimulants in respiratory depression from morphine. Fed. Proc. 3: 75, 1944.

Eight human subjects given 0.5 mg/kg of morphine SO<sub>4</sub> produced consistent respiratory depression which was maximal after one hour. Stimulants were given one hour after morphine. 0.1 - 0.4 mg/kg of amphetamine (s.c.) increased respiration back to normal. Metrazol and caffeine produced a weak stimulation but not as effective as amphetamine.

HARRIS, S. C. and F. J. FRIEND

Contribution of adrenals to morphine analgesia. Fed. Proc. 6: 124, 1947.

Using tail pinching method in rats after 5 mg/kg of morphine SO<sub>4</sub> (s.c.). In rats with bilateral adrenalectomy, but with cortices autotransplanted to the anterior chamber of the eye, the effect of morphine was reduced by 40%.

HERZ, A.

Über die Potenzierung der Morphinwirkung durch Spasmolytica vom Typ des Scopolamis. Naunyn-Schmiedebergs Arch. Exp. Path. Pharmak. 238: 110-111, 1960.

The cataleptic-narcotic action of morphine is enhanced by scopolamine, benactyzine, trihexyphenidyl, caramiphen and beperiden. Muscle tone is greatly reduced. The analgetic actions of morphine are enhanced some by scopolamine. A large enhancement of analgesia occurs with phenglutarimide and beperiden. The active drugs are considered to be acting in a similar fashion to chlorpromazine. Beperiden is 2 (Bicyclo [2:2:1] Hept-5-EN-2YL) phenyl-1-piperidenepropanol (it lowers Ach content of brain: see Haas, H. et al, Arch. Int. Pharmacodyn. 128: 204-252, 1960).

HURST, E. W. and O. L. DAVIS

Studies on the blood-brain barrier. III. Attempts to influence the passage of substances into the brain. Brit. J. Pharmacol. Chemotherap. 5: 147-164, 1950.

Using the reaction of rats to thermal stimuli the effect of a number of substances on morphine analgesia were studied by changes in mean reaction time. Of theocin, sodium lactate, hexamine, glycerol, adrenalin (.002 mg/100 g) + pituitrin and histamine (17.5 mg/100 g) only histamine and pituitrin + adrenaline changed morphine (0.2 mg/100 g) actions by potentiating analgesia. Also using various dyes it was found that neutral-red chloride (1 cc/100 g of 1%) given i.p. for four successive days prior to morphine produced a potentiation. No explanation was presented.

IVY, A. C., F. R. GOETZL and D. Y. BURRILL

Morphine-dextro-amphetamine analgesia. War Med. 6: 67-71, 1944.

21 healthy humans were given s. c. 16 mg of morphine  $\text{SO}_4$  and 20 mg dextroamphetamine. Recorded pain threshold by stimulating metal filling of tooth, critical flicker fusion, blood pressure, pulse rate and choice reaction time. D-amphetamine raised pain threshold of morphine about 60%. Mixture also raised CFF threshold above morphine, blood pressure, pulse rate were normalized and speeded choice RT. There was less nausea and vomiting following the mixture than with morphine alone. Drowsiness was much less with the mixture.



IVY, A. C., F. R. GOETZL, S. C. HARRIS and D. Y. BURRILL

The analgesic effect of intracarotid and intravenous injections of epinephrine in dogs and subcutaneous injections in man. Quart. Bull. Northwestern Univ. Med. Sch. 18: 298, 1944.

Using tooth pulp stimulation method in dogs they found that epinephrine, ephedrine and amphetamine all produced a potentiation of morphine analgesia. These drugs also had some analgesic action of their own. Similar studies on man confirmed the results.

KNOLL, J. and E. KOMLOS

Die Analgetische Wirkung des Atropins und sein Synergismus mit Morphin und Prostigmin. Acta Physiol. Hung. 2: 57-68, 1951.

Used mice on hot plate method and found that at a dose of between 50-100  $\mu$ g/kg atropine significantly potentiated the analgesic action of morphine at doses of 1-5 mg/kg. Similarly at a dose of 5 mg/kg scopolamine synergized the analgesic effects of morphine. Giving atropine plus prostigmine (0.1 mg/kg) plus morphine produced a degree of analgesia greater than any one or any combination of two of them.

KNOLL, J., E. KOMLOS and J. PORSZASZ

Analgesie und cholinesterasehemmung. Acta Physiol. (Budapest), 2: 478-491, 1951.

Used the mice on hot plate method to measure analgesic activity and cholinesterase activity of morphine, methadone, prostigmine, amidazophene, pethidine and atropine. No relationship was found between ChE activity and analgesia. A synergism of analgesic activity was found between morphine and prostigmine. Used 5 mg/kg of morphine and 0.1 mg/kg of prostigmine.

KOMLOS, E., J. PORSZASZ and J. KNOLL

Morphin-Prostigmin Synergismus. Acta Physiol. Acad. Sci. (Hung), 1: 77-90, 1950.

Used hot plate method in mice and found prostigmine to potentiate morphine at doses 0.8 - 5.0 of morphine and 0.1 of prostigmine.

There was also a potentiation of toxicity. Atropine (320 mg/kg) reduced the synergism of toxicity.

KULSRESHTHA, J. K. and P. N. SAXENA

Effect of atropine and hyoscine on morphine induced analgesia. Indian J. Physiol. Pharmacol. 1: 204-207, 1957.

No significant changes in the  $AD_{50}$  of morphine in rats was produced in combination with 150-300  $\mu$ g/kg or scopolamine HBr at 150  $\mu$ g/kg at 300  $\mu$ g/kg in rats scopolamine reduced analgesia.

MATSUMURA, M., S. TAKAORI and R. INOKI

Suppressive synergism of morphine with methamphetamine on the afferent pathways of the central nervous system in the cat. Jap. J. Pharmacol. 9: 67-74, 1959.

Cats under hexobarbital or spinal chord cut. Drugs given i.v. The effective dose of morphine was 6 mg/kg. Both morphine and methamphetamine suppressed cortical and intraspinal potentials of splanchnic afferent stimulation. Morphine was more potent at suppressing augmenting responses following repetitive stimulation of the medial lemniscus. Definite synergism was shown for the drugs on this augmenting response. Methamphetamine more strongly suppressed recruiting responses following repetitive stimulation of nucleus centre median. Possibly some synergism here. Morphine inhibited cortical responses but methamphetamine did not. No synergism shown in the cortex.

McKENZIE, J. S.

The effect of amiphenazole on morphine induced analgesia in mice. Aust. J. Exp. Biol. Med. Sci. 38: 47-59, 1960.

Used hot plate method in mice and found amiphenazole to potentiate morphine analgesia by i.p., s.c., oral routes. Effects on codeine were mild and variable. Suggest use in clinic to reduce necessary amount of narcotic.

MERCIER, F. and PAULETTE ETZENSPERGER

Potentialisation par la spartéine de l'actavite analgesique expérimentale. C. R. Soc. Biol. 148: 1431-1434, 1954.

Used the D'Amour and Smith method with the rat. Both temperature and time to reaction were measured. Sparteine (10 mg/kg) was given with:

morphine 1.5 mg/kg	potentiate
morphine 2.5 mg/kg	"
diamorphine 0.5 mg/kg	"
codeine 20.0 mg/kg	"
dihydrocodeine 10.0 mg/kg	antagonize
dihydrocodeine 10.0 mg/kg	no effect
dihydrone 1.0 mg/kg	"
piridosal 5.0 mg/kg	"
antipyrine 100.0 mg/kg	"
pyramiden 100.0 mg/kg	"
aspirine 60.0 mg/kg	potentiate
benzoxazolone 100.0 mg/kg	"

MERCIER, F. and PAULETTE MARINACCE

Influence exercée par un ester du diéthylaminoéthanol sur l'activité expérimentale de quelques analgésiques centraux. C. R. Soc. Biol. 145: 1340-1342, 1951.

Using hot plate method of Woolfe and Macdonald animals received either morphine (2.5 - 5.0 mg/kg), dihydrone (2.0 mg/kg) or pethidine (10 mg/kg) alone or in combination with 25 mg/kg of an ester of diethylaminoethanol (anisylhydrocinnamol), an antispasmodic agent. The degree of enhancement of analgesia was the same as if the dose of morphine had been increased by 100%.

MERCIER, F., J. MERCIER, PAULETTE ETZENSPERGER and D. ROUILLON

Reinforcement par les ganglioplégiques des activités analgésiques et anesthésique locale expérimentale. C. R. Soc. Biol. 148: 1448-1450, 1954.

Used rats with the method of Lespagnol and Mercier in which the intensity of hot plate heat to reaction is measured. 1.5 mg/kg of morphine was enhanced by Sparteine (10 mg/kg), pendiomide (2.5 mg/kg), penthonium (2.5 mg/kg), hexamethonium (2.5 mg/kg) and tetraethylammonium (2.5 mg/kg). Enhancement up to 100% for Sparteine and down to 3% for tetraethylammonium. The analgetic activity was not prolonged.

MILOSEVIC, M. P.

The analgesic activity of sympathomimetic amines. *Acta Med. Jugosl.* 12: 111-119, 1958.

Contact heat in mice showed weak action of adrenalin which is blocked by adrenergic blockers. Adrenergic blockers potentiated methadone. Nalorphine did not block epinephrine effect. Amphetamine and methamphetamine potentiated opiate. Adrenaline does not potentiate opiate.

MORRISON, J. L. and B. E. ABREU

Effects of d-, l-, and r-amphetamine on oxygen consumption of morphinized dogs. *Fed. Proc.* 2: 88, 1943.

O<sub>2</sub> consumption in dogs given 10 mg/kg morphine SO<sub>4</sub> plus 0.5 mg/kg of r-amphetamine, d-amphetamine, or l-amphetamine. R-amphetamine raised O<sub>2</sub> consumption 13% above basal morphine level, d-amphetamine 22% and l-amphetamine 15%.

NICKERSON, M.

Analgesic effects of nitrous oxide and meperidine alone and combined with amphetamine. *West. Soc. Clin. Rsch.* p. 541, April 1950.

Used human subjects with ice water immersion method and dolorimeter with meperidine (100 mg oral) N<sub>2</sub>O (33% by mask) with d-amphetamine (10-20 mg). Amphetamine potentiated the analgesia of meperidine, but not of N<sub>2</sub>O. Very little change under any condition with dolorimeter but good results with ice water method.

NICKERSON, M. and L. S. GOODMAN

Synergistic isonipecaïne-amphetamine analgesia. *Fed. Proc.* 6: 360-361, 1947.

Investigated human subjects with: cold water immersion, amperage applied to metal tooth filling and dolorimeter. 100 mg isonipecaïne + 10 mg amphetamine orally affected as a potentiation only on the "deep pain" of ice water technique. Considered a factor of more "subjective factor" of deep pain.

PORSZASZ, J., J. KNOLL and E. KOMLOS

Wirkung der parasymphomimetika auf die analgesia. *Acta Physiol. Acad. Sci. Hungaricae*, 2: 469-477, 1951.

Used a modified hot plate method with mice. Found a synergism of morphine, methadone, amidazophen, dolantin and hexalgon by: prostigmine, 0.1 mg/kg; physostigmine, 0.1 mg/kg; carbachol, 0.25 mg/kg. Of these carbachol gave greatest degree of effect. Tetraethylpyrophosphate at dose of 0.5 mg/kg gave no synergism. Concluded that effect is not due to parasymphomimetic action.

SADOVE, M. S., M. J. LEVIN, R. F. ROSE, L. SCHWARTZ and F. W. WITT

Chlorpromazine and narcotics in the management of pain of malignant lesions. *J.A.M.A.* pp. 626-628, June 1954.

Chlorpromazine in dose of 25 mg orally b.i.d. was given to 30 patients while reducing their narcotic dose by 50%. All patients maintained a satisfactory degree of analgesia and some were better on the mixture than previously on narcotics alone. Similarly nausea and vomiting were reduced.

SAXENA, P. N.

Mechanism of cholinergic potentiation of morphine analgesia. *Indian J. Med. Res.* 46: 653-658, 1958.

The experimenter used albino rats with bulldog clamp on tail as analgesic measurement technique. Drugs given alone and in combination with morphine. Pilocarpine, DFP, prostigmine and atropine tested at doses from 50 to 200  $\mu$ g/kg.

Drug	AD <sub>50</sub> in mg/kg
morphine	3.36
morphine + pilocarpine	2.13* (significant)
morphine + 100 DFP	3.45
morphine + 100 prostigmine + 200 $\mu$ g atropine	2.58

Thus atropine does not antagonize prostigmine potentiation of morphine. Also, DFP does not potentiate although it is a very strong anti ChE compound.

**SAXENA, P. N. and G. P. GUPTA**

Analgesia potentiating effects of ephedrine and methamphetamine.  
*J. Indian Med. Prof.* 4: 1553-1554 and 1599, 1957.

Analgetic level determined in rats by hot wire to tail method.

Drug	AD <sub>50</sub> of morphine (mg/kg)
morphine	4.29
phenylbutazone	lethal at these doses
morphine + ephedrine (6.25 mg/kg)	2.94
morphine + ephedrine (12.50 mg/kg)	2.17
morphine + methamphetamine (1.2 mg/kg)	3.58
morphine + methamphetamine (2.5 mg/kg)	3.12

**SCHAUMANN, W.**

Zur hypothese eines cholinergen Wirkungsmechanismus des Morphins. *Naunyn-Schmiedberg's Arch. Exp. Path. Pharmac.* 237: 229-240, 1959.

Using the Haffner method in mice it was found that morphine analgesia was potentiated by neostigmine and physostigmine. ChE inhibition by morphine previously found in vitro could not be confirmed in vivo. Irreversible peripheral inhibition of ChE had no effect on morphine analgesia. Morphine potentiated the toxicity of neostigmine, but not of physostigmine in mice. A cholinergic mechanism of morphine seems unlikely.

**SCHNEIDER, J. A.**

Reserpine antagonism of morphine analgesia in mice. *Proc. Soc. Exper. Biol. and Med.* 87: 614-615, 1954.

Using tail flick method in mice given 10.0 mg/kg of morphine SO<sub>4</sub> alone and with either reserpine (10.0 mg/kg) or chlorpromazine (10 mg/kg). Reserpine antagonized analgesia while chlorpromazine slightly enhanced it and considerably prolonged it.

**SCHNEIDER, J. A. and MARIE McARTHUR**

Potential action of ibogaine (Bogadin TM) on morphine analgesia. *Experimentia*, 12: 323-324, 1956.

Ibogaine, an indole alkaloid from *Tabernanthe iboga*, which has central stimulant properties was measured with the mouse tail flick method alone and in combination with 3 mg/kg morphine for analgesia. Alone, Ibogaine (6 - 24 mg/kg) had no effect but it significantly potentiated morphine analgesia by about 30 to 50%. It also increased LD<sub>50</sub> dose of morphine and Ibogaine LD<sub>50</sub> by five fold. It acted in a similar manner on codeine, methadone and ketobemidone. Aminopyrine was not potentiated.

**SIGG, E. B., G. CAPRIO and J. A. SCHNEIDER**

Synergism of amines and antagonism of reserpine to morphine analgesia. *Proc. Soc. Exper. Biol. and Med.* 97: 97-100, 1958.

Used tail flick in mice with s. c. doses: 5 mg/kg morphine SO<sub>4</sub> synergized

- |                                        |                       |
|----------------------------------------|-----------------------|
| 1. 5-hydroxytryptamine (5-10 mg/kg)    | Antagonized by        |
| 2. tryptamine (100 mg/kg)              | reserpine at 2.5      |
| 3. 5-hydroxytryptophane (30-400 mg/kg) | mg/kg but not by      |
| 4. amphetamine (3-5 mg/kg)             | isoreserpine, ipro-   |
| 5. mescaline (10-50 mg/kg)             | niazid (or iproniazid |
| 6. epinephrine (2-5 mg/kg)             | + reserpine).         |

Iproniazid, tryptophane or 5-hydroxyindoleacetic acid had no effect. Reserpine antagonism believed to be due to catechol amine depletion, but this has no effect on synergisms. Suggest retardation of demethylation as possible mechanism of synergism.

**SLAUGHTER, D. H.**

Neostigmine and opiate analgesia. *Arch. int. pharmacodyn.* 83: 143-148, 1950.

Used Wolff, Hardy and Goodell method on untrained subjects (6) with a double blind. Repeated measurements on same subjects used. Results in units of threshold over 150 min are by integration of difference from control: morphine SO<sub>4</sub> (16 mg), 150; morphine SO<sub>4</sub> (8 mg) + neostigmine methyl SO<sub>4</sub> (0.5 mg), 181; pantophon (20 mg), 91; pantophon (10 mg), 31; pantophon (10 mg) + neostigmine (0.5 mg), 57; dilaudid (6 mg), 327; dilaudid (3 mg),

156; dilaudid (3 mg) + neostigmine (0.5 mg), 223; codeine PO<sub>4</sub> (64 mg), 5; codeine (32 mg), -.7; codeine (32 mg) + neostigmine (0.5 mg), 51; neostigmine alone at 0.5 mg, 81. Neostigmine is said to reduce side effects of the opiates. Drugs were given I. M.

**SLAUGHTER, D. H. and D. W. MUNSELL**

Some new aspects of morphine action effects on pain. J. Pharmacol. exp. therap. 68: 104-112, 1940.

The investigators used responses of cats having their tail clamped (Eddy method). Drugs given s. c. atropine (.085 mg/kg) and prostigmine (.04 mg/kg) were given alone, in combination with morphine (1.0 mg/kg), and all three together. Prostigmine had no effect of itself but potentiated morphine about 100%. Atropine did not potentiate and reduced the effect of prostigmine potentiation.

**SZERB, J. C. and D. H. McCURDY**

Concentrations of morphine in blood and brain after intravenous injection of morphine in non-tolerant, tolerant and neostigmine-treated rats. J. Pharmacol. exp. therap. 118: 446-450, 1956.

Determining if neostigmine potentiates by increasing the passage of morphine into the brain by measuring brain morphine with and without neostigmine treatment. Rats were made tolerant by 37.5 mg/kg of morphine per day for seven days, 75.0 mg/kg for the next ten and 150 mg/kg for the last ten. Neostigmine by i. v. 0.2 mg/kg 20 min prior to morphine. Spontaneous activity and blood and brain morphine measured in tolerant and non-tolerant animals. Free morphine of blood and brain of tolerant rats was lower than non-tolerant. Neostigmine did not change the free or bound morphine in blood or brain, but did reduce tolerance to morphine and potentiate its action.

**TORDOS, L. and Z. JOBBAGYI**

Wirkung von Reserpin auf den Effekt der Analgetika. Acta Physiol. Acad. Sci. Hung. 13: 171-178, 1958.

Using a modified Woolfe-Macdonald method (hot plate) in rats reserpine (0.5 - 1.0 mg/kg) given 60 min prior, potentiated



morphine, pethidine and aminopyrine. Using tail flick method this potentiation was not found.

WIRTH, W.

Versuche zur Kombinierten Wirkung von Megaphen mit stark wirksamen Analgetica. Arch. f. exp. Path. u. Pharmacol. 222: 75-76, 1954.

Used rats and guinea pigs with a mechanical stimulation method. Found that the addition of chlorpromazine reduced threshold doses of morphine, meperidine and dormoran to one half to one third. Lethal dose of opiates was not changed.

WITKIN, L. B., C. F. HEUBNER, F. GALDI, E. O'KEEFE, P. SPITALETTA and A. J. PLUMMER

Pharmacology of 2-amino-indane HCl (SU-8629): A potent non-narcotic analgesic. J. Pharmacol. exp. therap. 133: 400-408, 1961.

Used tail flick method in mice and hot plate method with SU-8629 at 5 mg/kg and morphine at 2 mg/kg. Also used writhing from acetic acid, spontaneous motor activity, blood sugar, EST and gastric motility. Orally, SU-8629 is same potency as morphine but s.c. it is 1/5 as potent. SU-8629 is mild central stimulant similar to amphetamine but 1/4 as potent. SU-8629 not antagonized by nalorphine. Both SU-8629 and amphetamine potentiated morphine analgesia at same dose of 5 mg/kg but amphetamine was more potent.

WITKIN, L. B., M. MAGGIO and W. E. BARRETT

Independence of sedative and analgetic antagonizing effects of two reserpine esters. Proc. Soc. Exper. Biol. and Med. 101: 377-379, 1959.

The experimenters used tail flick in mice and found methylphenidate at 5 mg/kg to potentiate morphine analgesia by about 300%. Reserpine, syrosingopine and methyl-18-0 (dimethylamino benzoyl) reserpate (SU-5171) all antagonized morphine to about the same degree yet all three are not equipotent at sedation.

WITKIN, L. B., M. MAGGIO, E. O'KEEFE and F. GALDI

A study of some central nervous stimulants in mice. Fed.  
Proc. 19: 269, 1960.

The investigators used tail flick to heat in mice. Morphine analgesia was potentiated by methylphenidate and also by d-amphetamine. Metrazol was found to have no effect and reserpine antagonized analgetic action.

DISTRIBUTION LIST OF USAMHL REPORTS  
Project No. 6X60-01-001 Psychology Division

No. of  
Copies

AGENCY - DEFENSE

10 Armed Services Technical Information Agency, Arlington Hall Sta, Arlington 12, Virginia

AGENCY - US ARMY

- 1 Aberdeen Proving Ground, US Army Ordnance, Human Engineering Laboratory, Aberdeen Proving Ground, Maryland, Attn: Dr. J. D. Weiss, Director
- 1 Adjutant General, Department of the Army, Washington 25, D. C., Attn: AGTL
- 6 Army Attache, Box 79, Fleet Post Office, New York, N. Y., Attn: Col. John C. Cressler, Asst Army Attache (Medical)
- 1 Brooke Army Medical Center, US Army Surgical Research Unit, Library, Building 2653, Fort Sam Houston, Texas
- 1 Brooke Army Medical Center, Headquarters, Army Medical Service School, Physical Medicine Branch, Fort Sam Houston Texas, Attn: Captain Rachel Adams
- 1 Brooke Army Medical Service School, Publication Branch, Fort Sam Houston, Texas, Attn: Publication Branch
- 1 Brooke General Hospital, Medical Library, Box 151, Fort Sam Houston, Texas, Attn: Medical Librarian
- 1 Brooke General Hospital, Brooke Army Medical Center, Radioisotope Clinic, Fort Sam Houston, Texas
- 1 Chief Chemical Officer, Department of the Army, Washington 25, D. C., Attn: DCCa.O/SA
- 1 Chief of Ordnance, ORDTB, Washington 25, D. C., Attn: Res and Spec Proj Section
- 1 Chief of Research and Development, Chief Human Factors Research Division, Army Research Office, Room 3D-442, The Pentagon, Department of the Army, Washington 25, D. C.
- 1 Chief of Research and Development, Chief, Life Sciences Division, Room 3D-442, The Pentagon, Washington 25, D. C.
- 1 Chief of Research and Development, Army Research Office, Washington 25, D. C., Attn: Scientific Information Branch
- 1 Commanding General, US Army Alaska, APO 949, Seattle, Washington, Attn: Surgeon
- 1 Commanding General, US Army Europe, Medical Laboratory, Department of Microbiology, APO 180, US Forces, New York, N. Y.
- 1 Commanding General, US Army Europe, Medical Division, Plans and Operations Branch, APO 403, New York, N. Y., Attn: Chief Surgeon
- 1 Commanding General, US Army Hawaii, APO 957, San Francisco, California, Attn: Surgeon
- 1 Commanding General, US Army Japan, APO 343, San Francisco, California, Attn: Surgeon
- 1 Commanding General, US Army Ryukyu, Islands IX Corps, APO 331, San Francisco, California, Attn: Surgeon
- 1 Commanding General, Eighth US Army, APO 301, San Francisco, California, Attn: Surgeon
- 1 Commanding General, 7th Logistical Command, APO 612, San Francisco, California, Attn: Surgeon
- 1 Commanding General, I Corps Group, APO 358, San Francisco, California, Attn: Surgeon
- 1 Commander in Chief, US Army Pacific, APO 958, San Francisco, California, Attn: Surgeon
- 1 Fitzsimons General Hospital, Medical Technical Library, Denver 30, Colorado
- 1 Fitzsimons General Hospital, US Army Medical Research and Nutrition Laboratory, Denver, 30, Colorado
- 1 Picatinny Arsenal, Dover, New Jersey, Attn: ORDEB-VS3
- 2 Quartermaster Research and Engineering, Commanding General, Natick, Massachusetts, Attn: Technical Library
- 1 Quartermaster Research and Engineering Field Evaluation Agency, Fort Lee, Virginia, Attn: Technical Library
- 1 Redstone Arsenal, Commander, Redstone Arsenal, Alabama, Attn: ORDDW-HI

No. of  
Copies

US ARMY - CONTINUED

- 1 Surgeon General, Department of the Army, Professional Division, Washington 25, D. C.  
Attn: Psychiatry and Neurology Consultant
- 1 Valley Forge Army Hospital, Pheonixville, Pennsylvania, Attn: Medical Library
- 1 William Beaumont General Hospital, Medical Library, El Paso, Texas, Attn: Med. Librarian
- 1 US Army Air Defense, Human Research Unit, Fort Bliss, Texas, Attn: Library
- 1 US Army Armor Human Research Unit, Director, Fort Knox, Ky., Attn: Librarian
- 1 US Army Armor Infantry Human Research Unit, Director of Research, Post Office Box 2086,  
Fort Benning, Georgia, Attn: Library
- 1 US Army Chemical Research and Development Laboratories, Technical Library, Building 330,  
Army Chemical Center, Maryland, Attn: Librarian
- 2 US Army Medical Center, Director of Medical Research ORDC, Army Chemical Center, Maryland
- 1 US Army Chemical Corps, Biological Laboratories, Fort Detrick, Maryland, Attn:  
Documents Librarian
- 1 US Army Dispensary, Springfield Armory, Springfield, Massachusetts
- 1 US Army Engineer Research and Development Laboratories, Fort Belvoir, Virginia, Attn:  
Technical Documents Center
- 1 US Army Environmental Hygiene Agency, Commanding Officer, Army Chemical Center, Maryland
- 1 US Army Hospital, Fort Lee, Virginia, Attn: Medical Library
- 1 US Army Liaison Group, Project Michigan, Willow Run Laboratories, Ypsilanti, Michigan,  
Attn: Lt. Col. R. S. Johnston, Executive Officer
- 1 US Army Medical Command Japan, Medical General Laboratory (406) APO 343, San Francisco,  
California, Attn: Colonel Carl F. Tessner, MC, Commanding
- 1 US Army Leadership, Human Research Unit, Library, Post Office Box 787, Presidio of  
Monterey, California
- 2 US Army Medical Liaison Branch, Office of the Chief Surgeon, Corgas Hospital, Balboa  
Heights, Canal Zone
- 3 US Army Medical Research and Development Command, Commanding General, Main Navy Bldg.,  
Washington 25, D. C.
- 1 US Army Medical Research Unit, Commanding Officer, Panama Field, Fort Clayton, Canal Zone
- 1 US Army Medical Unit, Commanding Officer, Fort Detrick, Maryland
- 1 US Army Medical Research Unit, Institute for Medical Research, Kuala Lumpur, Malaya
- 1 US Army Ordnance Arsenal, Frankford, Philadelphia 37, Pennsylvania, Attn: ORDBA-1734/65-1  
Mr. A. Charles Karr
- 2 US Army Ordnance Tank-Automotive Command, Detroit Arsenal, Center Line, Michigan
- 1 US Army Medical Research and Development Command, Office of The Surgeon General, Attn:  
MEDDH-HR
- 1 US Army Standardization Senior Representative, US Army Standardization Group, Canada  
c/o US Army Attache, US Embassy, Ottawa, Ontario, Canada, Attn: Col Joseph R. Blair
- 1 US Army Tropical Research Medical Laboratory, APO 851, New York, N. Y.
- 1 US Army Research Office, OORD, Department of the Army, Washington 25, D. C., Attn:  
Scientific Info. Branch
- 1 US Atomic Energy Commission, Division of Biology and Medicine, Washington 25, D. C.  
Attn: Robert L. Corsbie
- 3 Walter Reed Army Institute of Research, Walter Reed Army Medical Center, Director,  
Washington 12, D. C.
- 1 Walter Reed Army Institute of Research, Division of Neuropsychiatry, Office of the Dir.,  
Washington 12, D. C.
- 1 Walter Reed Army Hospital, Army Audiology and Speech Center, Forest Glen Section,  
Washington 12, D. C.
- 1 Walter Reed Army Medical Center, Commanding Officer, US Army Prosthetics Research  
Laboratory, Washington 12, D. C.
- 1 Walter Reed Army Medical Center, US Army Medical Service, Historical Unit, Washington 12,  
D. C., Attn: Mrs. Esther E. Rohlader, Asst Chief, General Reference and Research Br

# AGENCY - US NAVY

No. of  
Copies

- 2 Bureau of Medicine and Surgery, Director, Research Division, Department of the Navy, Washington 25, D. C.
- 2 Bureau of Naval Weapons (PLI-31) Department of the Navy, Washington 25, D. C.
- 1 Bureau of Naval Weapons (PAAE-13), Department of the Navy, Washington 25, D. C., Attn: Capt. Walton L. Jones, (SC) USN
- 2 Bureau of Yards and Docks, Department of the Navy, Washington 25, D. C., Attn: Code D-440
- 1 Chief of Naval Air Technical Training, US Naval Air Station, Memphis 75, Tennessee, Attn: Staff Medical Officer
- 1 Naval Air Crew Equipment Laboratory, Naval Air Material Center, Philadelphia 12, Pa.
- 1 Naval Air Reserve Training, Staff Medical Officer, US Naval Air Station, Glenview, Ill.
- 1 Naval Medical Research Institute, Technical Reference Library, National Naval Medical Center, Bethesda 14, Maryland
- 1 Naval Medical Research Laboratory, Box 100, Naval Submarine Base, New London, Conn.
- 1 Office of Naval Research, (Code 454) Department of the Navy, Washington 25, D. C.
- 1 Office of Naval Research (Code 408) Department of Navy, Special Assistant for Medical and Allied Sciences, Washington 25, D. C.
- 10 Office of Naval Research Branch Office, Navy 100, Box 39, Fleet Post Office, New York, New York, Attn: Victoria S. Greswold, Head, Administrative and Tech Service Branch
- 1 US Naval Air Development Center, Aeronautical Instrument Laboratory, Johnsville, Pa. Attn: Simulation Branch
- 1 US Naval Air Development Center, Aviation Medical Acceleration Laboratory, Johnsville, Pennsylvania, Attn: Library
- 1 US Naval Civil Engineering Laboratory, Port Hueneme, California, Attn: Code 313
- 1 US Naval Medical Neuropsychiatric Research Unit, San Diego 52, California
- 2 US Naval Missile Center, Point Mugu, California, Attn: Technical Library
- 1 US Naval Medical School, National Naval Medical Center, Commanding Officer, Bethesda 14, Maryland
- 1 US Naval Ordnance Test Station, The Medical Officer (Code 88), Station Hospital, China Lake, California
- 1 US Naval Radiological Defense Laboratory, Commanding Officer and Director (222), San Francisco 24, California
- 1 US Naval Research Laboratory, Code 2027, Washington 25, D. C.
- 1 US Naval School of Aviation Medicine, Aviation Psychology Laboratory, US Naval Aviation Medical Center-54, Pensacola, Florida
- 2 US Naval School of Aviation Medicine, Director of Research, US Naval Aviation Center 54, Pensacola, Florida
- 2 US Naval Training Device Center, Human Engineering Department, Code 34, Port Washington, New York
- 1 US Naval Weapons Plant, Experimental Diving Unit, Washington 25 D. C.

## AGENCY - AIR FORCE

- 1 Air Force Flight Test Center, Human Factors Branch (FTFHD) Edwards Air Force Base, California, Attn: Chief, of Human Factors
- AIR RESEARCH AND DEVELOPMENT COMMAND
- 2 Air Research and Development Command, US Air Force, (RDGRL) Andrews Air Force Base, Washington 25, D. C.
  - 1 Air Research and Development Command, Office of Scientific Research, Behavioral Sciences Division, Washington 25, D. C.
  - 5 Air Research and Development Command, Operations Office, Laurence G. Hanscom Field, Bedford, Massachusetts, Attn: CORHA
  - 3 Air Research and Development Command, Operational Applications Office, (OORH), Laurence G. Hanscom Field, Bedford, Massachusetts
  - 1 Air Research and Development Command, Wright Air Development Center, US Air Force, Wright-Patterson Air Force Base, Ohio, Attn: D. G. Pitts, Vision

No. of  
Copies

AIR FORCE - CONTINUED

AIR RESEARCH AND DEVELOPMENT COMMAND CONTINUED

- 1 Wright Air Research and Development Command, (WERDAS Aerospace Medical Library) YADD  
Wright-Patterson Air Force Base, Ohio
- 1 Wright Air Development Division, US Air Force, Aerospace Medical Division, Wright-  
Patterson Air Force Base, Ohio. Attn: WERDAS, Lt Col. Stanley L. Pleha
- 1 Wright Air Development Division, Wright-Patterson Air Force Base Ohio, Attn: WERDAS
- 2 Air Training Command, Office of The Surgeon General-P, Randolph Air Force Base, Texas
- 1 Arctic Aeromedical Laboratory, APO 731, Seattle, Washington, Attn: Library
- 1 Assistant for Ground Safety, DCS/P, US Air Force, Washington 25, D. C.
- 1 Ent Air Force Base, Command Surgeon, US Army Air Defense Command, Colorado, Springs,  
Colorado, Attn: Colonel S. J. Newson, MC
- 1 Ent Air Force Base, Commander Air Development Center, OMO, Human Factors, Box 15,  
Colorado Springs, Colorado
- 1 Human Factors Division, Director of Research and Technology, US Air Force, Washington  
25, D. C., Attn: Chief, Human Factors Division
- 3 Langley Research Center, National Aeronautics and Space Administration, Langley Field,  
Virginia, Attn: Librarian
- 1 Rome Air Development Center, Air Research and Development Command, US Air Force, Griffiss,  
Air Force Base, New York
- 2 US Air Force Aerospace Medical Center, SAM, (SAMSCHE-P) Brooks Air Force Base, Texas
- 1 US Air Force Hospital, US Air Force Aerospace Medical Center (ATC), Lackland Air Force  
Base, Texas, Attn: Medical Librarian
- 1 US Air Force, Strategic Air Command, (SUP3), Offutt Air Force Base, Nebraska

GOVERNMENTAL - AGENCIES

- 1 Argonne National Laboratory, 9700 South Cass Avenue, Argonne, Illinois, Attn: Hoylande  
D. Young
- 1 Central Intelligence Agency, 2430 E. Street, N.W., Washington, D. C., Attn: 1331, RES Bldg.
- 1 Civil Aeromedical Research Institute, Federal Aviation Agency, Post Office Box 1082,  
Oklahoma City, Oklahoma
- 2 Library of Congress, Science and Technology Division, Washington 25, D. C., Attn: Dr.  
A. J. Jacobus
- 1 National Institute of Health, Library, Building 10, Room 3N118, Bethesda 14, Maryland,  
Attn: Acquisitions Section
- 1 National Library of Medicine, 7th Street & Independence Ave, S.W., Washington 25, D. C.
- 1 National Institutes of Health, Division of Research Grants, Information Office, Bethesda  
14, Maryland
- 1 Prosthetic and Sensory Aids Service, Chief, Research and Development Division, Veterans  
Administration, 252 7th Ave., New York 1, New York, Attn: Eugene F. Murphy, Ph.D.

OTHER AGENCIES

- 1 Arctic Health Research Center, Library, Post Office Box 960, Anchorage, Alaska
- 1 Bell Telephone Laboratories, Murray Hill, New Jersey, Attn: Max S. Schoeffler
- 1 Boeing Airplane Company, Central Medical Library, Box 11-40, Post Office Box 3707,  
Seattle, Washington
- 1 Boeing Airplane Company, Chief, Space Medicine Section, Aero-Space Division, Seattle 24,  
Washington, Attn: Dr. Romney H. Lowry, (Box 19-29)
- 1 Boeing Airplane Company, Library, Wichita Division, Wichita 1, Kansas
- 1 Chrysler Corporation, Box 1118, Engineering Research Department 921, Detroit 31, Michigan,  
Attn: John Versace, Engr. Psychologist
- 1 Electric Boat Division, General Dynamics Corp., Human Factors Section, Groton, Conn
- 1 Ford Motor Company, Technical Information Section, Scientific Laboratory, Post Office  
Box 2053, Dearborn, Michigan
- 1 General Electric Company, Technical Military Planning Operation, Library, 735 State  
Street, Santa Barbara, California

No. of  
Copies

OTHER AGENCIES - CONTINUED

- 1 General Electric Company, Advanced Electronics Center, at Cornell University, Ithaca, New York
- 1 IBM Research Center, Biophysics Group, Box 218, Yorktown Heights, New York, Attn: Dr. W. R. Uttal
- 1 ITT Laboratories, Human Factors Division, 390 Washington Avenue, Nutley 10, New Jersey, Attn: Richard G. Lazar, Human Factors Analyst
- 1 John Crerar Library, 85 East Randolph Street, Chicago 1, Illinois
- 1 Mayo Clinic, Section of Biophysics, Rochester, Minnesota, Attn: Dr. Kenneth N. Ogle
- 1 Mercy Hospital, Anesthesia Research Laboratory, Locust and Pride Streets, Pittsburgh 19, Pennsylvania
- 1 Motorola Incorporated, Systems Research Laboratory, 9330 Indiana Avenue, Riverside, California, Attn: R. E. Freese, Tech Info Supervisor
- 7 National Aeronautics and Space Administration, 1520 H Street, N. W., Washington 25, D. C., Attn: Bertram A. Mulcahy, Asst Director for Technical Information
- 2 Rheem Califorme Corporation, 1020 North La Brea, Los Angeles 38, California, Attn: Mr. W. R. Deutsch, Head, Research and Development
- 1 Space Technology Laboratory, Subcommittee on Noise, 327 South Alvarado Street, Los Angeles 57, California
- 1 Syceas Research Center, Lockheed Electronics Company, Bedminster, New Jersey, Attn: R. E. Wever, Box 37
- 1 Yerkes Laboratories of Primate Biology, Incorporated, Orange Park, Florida

MEDICAL COLLEGE/SCHOOL LIBRARIES AND DEPARTMENTS

- 1 Albany Medical College Library, New Scotland Avenue, Albany 8, New York
- 1 Antioch College, Engineering Psychology Research Project, Yellow Springs, Ohio
- 1 Bowman Gray School of Medicine Library, Winston-Salem, North Carolina
- 1 Brown University, Department of Psychology, Providence 12, Rhode Island, Attn: Prof. L. A. Riggs
- 1 Brown University, Department of Psychology, Providence 12, Rhode Island, Attn: Prof. Harold Schlosberg, Consultant
- 1 College of Medical Evangelists, White Memorial Medical Library, 1720 Brooklyn Avenue, Los Angeles 33, California
- 1 College of Medical Evangelists, Vernier Radcliffe Memorial Library, Loma Linda, California
- 1 College of Physicians of Philadelphia, Library, 19 South 22nd Street, Philadelphia, 3, Pennsylvania
- 1 Columbia University, Department of Psychology, New York 27, New York, Attn: Dr. C. H. Graham
- 1 Columbia University Medical Library, 630 West 168th Street, New York 32, New York
- 1 Cornell University Medical College Library, 1300 York Avenue, New York 21, New York
- 1 Creighton University, Medical-Pharmacy Library, 1401 Davenport, Omaha 2, Nebraska
- 1 Emory University, Department of Psychology, Atlanta 22, Georgia, Attn: Dr. E. A. Alluisi
- 1 Florida State University, Department of Psychology, Tallahassee, Florida, Attn: Dr. W. F. Dawson
- 1 Fordham University, Fordham Road, New York 58, New York, Attn: R. T. Zegers, Sr.
- 2 George Washington University, Human Resources Research Office, P. O. Box 3596, Washington 7, D. C., Attn: Library
- 1 Harvard School of Public Health, 1 Shattuck Street, Boston 15, Massachusetts, Attn: Dr. Albert Damon
- 1 Harvard Medical Library, 25 Shattuck Street, Boston 15, Massachusetts, Attn: Library
- 1 Indiana University, Department of Psychology, Bloomington, Indiana, Attn: Prof. R. C. Davis
- 1 Indiana University Medical Center, School of Medicine Library, 1100 West Michigan Street, Indianapolis 7, Indiana
- 1 Indiana University Medical Center, 1199 West Michigan Street, Indianapolis 7, Indiana, Attn: Dr. Harris B. Shumacker Jr., Prof. of Surgery

No. of  
Copies

MEDICAL COLLEGE/SCHOOL LIBRARIES AND DEPARTMENTS

- 1 Indiana University, Anatomy-Physiology Department, Bloomington, Indiana, Attn: Dr. Sid Robinson
- 1 Jefferson Medical College Library, 1025 Walnut Street, Philadelphia 7, Pennsylvania
- 1 John Hopkins University, Welch Medical Library, 1900 E. Monument Street, Baltimore 5, Maryland
- 1 Kansas State University, Psychology Division, Manhattan, Kansas, Attn: Dr. William Bevan
- 1 Marquette University, Medical-Dental Library, 560 North 16th Street, Milwaukee, Wisconsin
- 1 Medical College of Virginia, Tompkins-McCaw Library, Richmond 19, Virginia
- 1 New York University, College of Engineering Research Division, 252 7th Avenue, New York New York
- 1 New York Academy of Medicine Library, 2 East 103rd Street, New York 29, New York
- 1 New York University Medical Center, Medical Library, 550 First Avenue, New York 16, N. Y.
- 1 Northwestern University, Department of Psychology, Evanston, Illinois, Attn: William A. Hunt
- 1 Northwestern University Medical School, Archibald Church Library, 303 E Chicago Avenue, Chicago 11, Illinois
- 1 Ohio State University Research Center, Director, Aviation Psychology Laboratory, 1314 Kennear Road, Columbus 12, Ohio
- 1 Ohio State University Research Center, Psycholinguistics Laboratory, 1314 Kennear Road, Columbus 12, Ohio
- 1 Ohio State University, Topaz Library, School of Optometry, 338 West 10th Avenue, Columbus 10, Ohio
- 1 Ohio State University, School of Optometry, Columbus 10, Ohio, Attn: Glenn A. Fry, Dir.
- 1 Rush Medical College Library, 1758 West Harrison Street, Chicago 12, Illinois, Attn: Librarian
- 1 Stanford University, Lane Medical Library, 300 Pasteur Road, Palo Alto, California
- 1 Stanford University, Department of Physiology, Stanford, California, Attn: J. M. Crismon, M.D.
- 1 State University of Iowa, College of Medicine Library, Medical Laboratories Building, Iowa City, Iowa
- 1 State University of New York, Downstate Medical Center, Department of Anesthesiology, 459 Clarkson Avenue, Brooklyn 3, New York
- 1 State University of New York, Downstate Medical Center, Medical Library, 450 Clarkson Avenue, Brooklyn 3, New York
- 1 Texas Medical Center Library, Jesse H Jones Library Building, Houston 25, Texas
- 2 Tufts University Institute for Applied Experimental Psychology, Medford, Massachusetts
- 1 Tulane University, School of Medicine, 1430 Tulane Avenue, New Orleans 12, Louisiana, Attn: G. E. Burch, M.D., Professor of Medicine
- 1 West Virginia University, Medical Center Library, Morgantown, West Virginia
- 1 University of Alabama, Medical Center Library, 1919 Seventh Avenue South, Birmingham 3, Alabama
- 1 University of Arkansas, Medical Center Library, 4301 West Markham, Little Rock, Arkansas
- 1 University of Buffalo, Department of Psychology, Buffalo 14, New York
- 1 University of Buffalo, Health Sciences Library, Copen Hall 141, Buffalo 14, New York, Attn: Librarian
- 1 University of California, Medical Center, Biomedical Library, Los Angeles 24, California
- 1 University of Chicago, Laboratory of Physiological Psychology, Faculty Exchange Box 184, Chicago 37, Illinois, Attn: W. D. Neff
- 1 University of Illinois Airport, Aviation Psychology Laboratory, Savoy, Illinois
- 1 University of Illinois, Department of Psychology, Urbana, Illinois, Attn: W. E. Kappauf
- 1 University of Illinois, Department of Psychology, Training Research Laboratory, 45 Lincoln Hall, Urbana, Illinois, Attn: Professor Lawrence M Stolorow, Ph.D.
- 1 University of Illinois, Documents Division, Urbana, Illinois
- 1 University of Illinois, Speech Hearing Clinics, 601 E. John Street, Champaign, Illinois, Attn: John J O'Neill
- 1 University of Kansas, Medical Center, Clendening Medical Library, Kansas City, Kansas
- 3 University of Louisville, School of Medicine Library, 101 W. Chestnut, St. Louisville, Ky.



No. of  
Copies

MEDICAL COLLEGE/SCHOOL LIBRARIES AND DEPARTMENTS

- 1 University of Miami, School of Medicine, Department of Ophthalmology, Miami 36, Florida
- 1 University of Michigan, 4506 Kresge Medical Research Building, Ann Arbor, Michigan.  
Attn: Dr. Kerle Lawrence
- 1 University of Michigan, Serials and Documents Section, General Library, Ann Arbor, Michigan
- 1 University of Minnesota Library, Serials Division, Minneapolis 14, Minnesota
- 1 University of Nebraska, College of Medicine Library, 42nd and Daway Avenue, Omaha, Nebraska, Attn: Librarian
- 1 University of North Carolina, Division of Health Affairs Library, North Carolina Memorial Hospital, Chapel Hill, North Carolina
- 1 University of North Carolina, Department of Medicine, Chapel Hill, North Carolina.  
Attn: Dr. Richard L. Dobson
- 1 University of North Carolina, School of Medicine, Department of Physiology, Chapel Hill, North Carolina
- 1 University of Oklahoma, Medical Center Library, 801 N. E. 13th Street, Oklahoma City 4, Oklahoma
- 1 University of Oregon, Medical School Library, Portland 1, Oregon, Attn: Librarian
- 1 University of Pittsburgh, Falk Library of the Health Professions, Pittsburgh 13, Pa
- 1 University of Pittsburgh, Graduate School of Public Health, Pittsburgh 13, Pa
- 1 University of Rochester, Department of Psychology, Rochester 20, New York, Attn: Dr. S. D. S. Spragg
- 2 University of Tennessee, College of Medicine, Clinical Physiology - Institute of Clinical Investigations, 62 South Dunlap, Memphis 3, Tennessee
- 1 University of Tennessee, Medical Units, Mooney Memorial Library, 62 South Dunlap, Memphis 3, Tennessee
- 1 University of Utah, Medical Library, Salt Lake City, 12, Utah
- 1 University of Vermont, College of Medicine Library, Burlington, Vermont
- 1 University of Virginia, Division of Educational Research, Old Cabell Hall, Charlottesville, Virginia
- 1 University of Virginia, Medical Library, University Hospital, Charlottesville, Virginia
- 1 University of Virginia, Psychological Laboratory, Peabody Hall, Charlottesville, Virginia
- 1 University of Washington, Health Sciences Library, Seattle 5, Washington
- 1 University of Wisconsin, Psychological Abstracts, 600 N. Park St., Madison 6, Wisconsin
- 1 University of Wisconsin, Medical School Library, S.M.I. Building, N Charter Street, Madison 6 Wisconsin
- 1 University of Wisconsin, Physiology Studies Division, 600 N. Park Street, Madison 6, Wisconsin, Attn: Janice Gans, Administrative Assistant

FOREIGN

- 1 American Embassy, Office of the United States Army Attache, Foreign Service of the United States of America, London, England, Army Attache, (Medical)
- 1 British Army Medical Liaison Officer, British Army Staff, British Embassy, Washington 8, D. C.
- 1 British Navy Staff Office, Benjamin Franklin Station, Post Office Box 165, Washington, D. C., Attn: Surgeon, Captain F. P. Ellis
- 10 Canadian Liaison Office, US Army, Office of The Surgeon General, Room 1709A, Main Navy Building, Washington 25, D. C.
- 2 Defence Research Member, Canadian Joint Staff, 1450 Massachusetts Avenue, N. W., Washington 8, D. C.
- 1 J. J. Thomson Physical Laboratory, Reading University, Whiteknights Park, Reading, England  
Attn: Professor R. W. Ditchburn
- 1 Karolinska Institute, Department of Histology, Stockholm 60, Sweden, Attn: Dr. Jan Wersall
- 1 Royal Society of Medicine, Library, 1 Wimpole Street, London W.1, England
- 1 Università Di Pisa, Istituto De Fisiologia, Pisa, Italy, Attn: Professor Giuseppe Moruzzi

No. of  
Copies

FOREIGN

- 1 Institute of Occupational Health, Hartmaninkatu 1, Helsinki, Finland, Attn: Dr. Martti J. Karvonen, Director of the Physiological Department
- 2 University of Western Ontario, Medical School, Department of Biophysics, South Street, London, Ontario, Canada, Attn: Dr. Allen C. Burton
- 1 University of Western Ontario, Department of Physiology, London, Ontario, Canada, Attn: Professor J. A. F. Stevenson

AD Accession No.

US Army Medical Research Lab., Ft. Knox, Ky.  
THE SYNERGISM OF AUTONOMIC DRUGS ON OPIATE  
OR OPIOID-INDUCED ANALGESIA: A DISCUSSION  
OF ITS POTENTIAL UTILITY AND AN ANNOTATED  
BIBLIOGRAPHY - W. O. Evans

Report No. 554, 10 Oct 62, 26 pp & ii -  
Project No. 6X60-01-001, Unclassified Report  
A diverse group of autonomically active drugs has been shown to potentiate an opiate-induced analgesia. Of these drugs, a d-amphetamine-opiate mixture offers an increase in analgetic effects of 60 to 100%, while minimizing the side effects of the opiate. The potential of this combination for use when traumatic injury occurs in situations in which shock is a particular difficulty is discussed. An annotated bibliography is presented on the potentiation of opiate-induced analgesia by autonomic drugs.

UNCLASSIFIED

1. Psychopharmacology
2. Analgesia
3. Opiate Potentiation

AD Accession No.

US Army Medical Research Lab., Ft. Knox, Ky.  
THE SYNERGISM OF AUTONOMIC DRUGS ON OPIATE  
OR OPIOID-INDUCED ANALGESIA: A DISCUSSION  
OF ITS POTENTIAL UTILITY AND AN ANNOTATED  
BIBLIOGRAPHY - W. O. Evans

Report No. 554, 10 Oct 62, 26 pp & ii -  
Project No. 6X60-01-001, Unclassified Report  
A diverse group of autonomically active drugs has been shown to potentiate an opiate-induced analgesia. Of these drugs, a d-amphetamine-opiate mixture offers an increase in analgetic effects of 60 to 100%, while minimizing the side effects of the opiate. The potential of this combination for use when traumatic injury occurs in situations in which shock is a particular difficulty is discussed. An annotated bibliography is presented on the potentiation of opiate-induced analgesia by autonomic drugs.

UNCLASSIFIED

1. Psychopharmacology
2. Analgesia
3. Opiate Potentiation

AD Accession No.

US Army Medical Research Lab., Ft. Knox, Ky.  
THE SYNERGISM OF AUTONOMIC DRUGS ON OPIATE  
OR OPIOID-INDUCED ANALGESIA: A DISCUSSION  
OF ITS POTENTIAL UTILITY AND AN ANNOTATED  
BIBLIOGRAPHY - W. O. Evans

Report No. 554, 10 Oct 62, 26 pp & ii -  
Project No. 6X60-01-001, Unclassified Report  
A diverse group of autonomically active drugs has been shown to potentiate an opiate-induced analgesia. Of these drugs, a d-amphetamine-opiate mixture offers an increase in analgetic effects of 60 to 100%, while minimizing the side effects of the opiate. The potential of this combination for use when traumatic injury occurs in situations in which shock is a particular difficulty is discussed. An annotated bibliography is presented on the potentiation of opiate-induced analgesia by autonomic drugs.

UNCLASSIFIED

1. Psychopharmacology
2. Analgesia
3. Opiate Potentiation

AD Accession No.

US Army Medical Research Lab., Ft. Knox, Ky.  
THE SYNERGISM OF AUTONOMIC DRUGS ON OPIATE  
OR OPIOID-INDUCED ANALGESIA: A DISCUSSION  
OF ITS POTENTIAL UTILITY AND AN ANNOTATED  
BIBLIOGRAPHY - W. O. Evans

Report No. 554, 10 Oct 62, 26 pp & ii -  
Project No. 6X60-01-001, Unclassified Report  
A diverse group of autonomically active drugs has been shown to potentiate an opiate-induced analgesia. Of these drugs, a d-amphetamine-opiate mixture offers an increase in analgetic effects of 60 to 100%, while minimizing the side effects of the opiate. The potential of this combination for use when traumatic injury occurs in situations in which shock is a particular difficulty is discussed. An annotated bibliography is presented on the potentiation of opiate-induced analgesia by autonomic drugs.

UNCLASSIFIED

1. Psychopharmacology
2. Analgesia
3. Opiate Potentiation